

Total Joint Repair Complications

Inpatient Physical Therapy Inservice

Nathaniel Yomogida

2024-09-25

Table of contents

1	Total Joint Arthroplasty Problems	2
2	Causes of failed discharge	2
3	Orthostatic Hypotension & Intolerance	2
3.1	Orthostatic Hypotension (OH)	2
3.2	Orthostatic Intolerance (OI)	3
3.3	Patient Presentation	3
3.4	Etiology	3
3.5	Pathophysiology	3
3.6	Pre-op Risk factors	4
3.7	Predictors	4
3.8	Pharmacological management	4
3.9	Hydration	4
3.10	Caffeine / Caffeine withdrawal	5
3.11	Sitting up	5
3.12	Exercise	5
3.13	Patient Education	5
3.14	Inpatient approach	6
4	Dropfoot	6
4.1	Etiology	6
4.1.1	THA	6
4.1.2	TKA	7
4.2	Epidemiology	7
4.2.1	THA	7
4.2.2	TKA	7
4.3	Common fibular nerve vs Tibial Nerve	7
4.4	Patient Presentation	7
4.4.1	Sensory symptoms	7

4.4.2	Motor Symptoms	7
4.4.3	Onsert	8
4.5	Management	8
4.5.1	Patient education	8
4.5.2	Positioning	8
4.5.3	Stretching	8
5	Anemia	8
5.1	Definition	8
5.2	Secondary complications	8
5.3	Management	8
	References	9

1 Total Joint Arthroplasty Problems

Post-operative complications can cause inpatient physical therapists to attempt therapy on a patient multiple times or even prevent a patient's discharge, which causes increased load on the physical therapists and PTAs, physical therapy administration, nursing, orthopedic department, and other aspects of the hospital.

2 Causes of failed discharge

- Orthostatic intolerance¹
- Insufficient muscle strength
- Poor sensation

3 Orthostatic Hypotension & Intolerance

3.1 Orthostatic Hypotension (OH)

[Orthostatic hypotension](#) refers to the clinical decrease in blood pressure associated with changes in position.

Table 1: Types of Orthostatic Hypotension

Type	Definition
Classic	Decrease in SBP of >20mmHg or DBP 10mmHg after 3 minutes of standing or head-up tilt (HUT) 60° on a tilt table ²
Initial	Temporary BP decrease of >40mmHg SBP or >20mmHg DBP within 15 seconds of standing ² .

3.2 Orthostatic Intolerance (OI)

Orthostatic intolerance is a presentation of symptoms associated with a sitting or standing position including:

- Dizziness¹
- Nausea¹
- Vomiting¹
- Blurred vision¹
- Syncope¹

3.3 Patient Presentation

On average, most orthostatic events occur within the first 12 hours *after* the surgical procedure, but can occur up to 48 hours after surgery¹.

3.4 Etiology

The causes of post-op orthostatic hypotension include:

- Surgical stress response¹
- Pain-induced¹
- Post-op Opioid administration¹
- Residual effects of Anesthesia¹
- Hypovolemia¹
- Acute anemia¹
- Preexisting Orthostatic Intolerance¹

3.5 Pathophysiology

Although there the pathophysiologic mechanism of orthostatic intolerance is not fully understood, there is a widely accepted theorized mechanism of orthostatic intolerance.

1. Standing up leads to a decrease in blood pressure rostrally and an increase in BP caudally.
2. The blood shifts below the diaphragm to the venous capacitance system³.
3. The fluid shift causes a decrease in venous return, ventricular filling, cardiac output, and blood pressure³.
4. This gravity-induced BP change is sensed by arterial baroreceptors in the aortic arch and carotid sinus³.

-
1. The body dysfunctionally has a diminished vasopressor response and absent baroreflex to these pressure changes¹.
 2. Due to the diminished autonomic response, the rostral blood pressure decreases.
 3. Decreased rostral blood pressure results in cerebral hypoperfusion¹.

4. Cerebral hypoperfusion can result in the syndrome of symptoms known as orthostatic intolerance¹.

3.6 Pre-op Risk factors

Patients in these groups had statistically significantly higher rates of OI:

Pre-op factors

- Older age¹
- Female¹
- THA > TKA or UKA¹
- Non-recreational drug users¹
- Lower preoperative diastolic BP¹

Perioperative factors

- Spinal +/- monitored anesthesia care > General +/- spinal¹
- Tramadol use
- No oxycodone use
- Increased PACU IVF
- Lower PACU Hgb

3.7 Predictors

When the above differences were examined using a multivariable analysis, only 4 items were found to significantly impact the odds of having orthostatic intolerance:

- Female gender (4.19 OR)¹
- THA surgery (vs TKA) (4.86 OR)¹
- Spinal + MAC anesthesia 2.35 OR (compared to spinal + general)¹.
- Bupivacaine spinal medication 1.79 OR (compared to Ropivacaine)¹.

3.8 Pharmacological management

There are pharmacologic measures that reduce orthostatic hypotension. The problem is that pharmacologic interventions that improve OH cause other cardiac side effects, primarily supine hypertension and ventricular hypertrophy⁴. In addition, since physical therapists cannot prescribe medications this is irrelevant to the profession.

3.9 Hydration

A systematic review by Figueroa³ found that drinking 16-oz of cold water can improve OH and related symptoms by expanding the plasma volume³. Within a few minutes, the cold water produces a *pressor effect*, which results in improve [orthostatic hypotension](#) by increasing SBP by >20 mmHg for ~2 hours and reducing symptoms of [orthostatic intolerance](#)³.

3.10 Caffeine / Caffeine withdrawal

According to a systematic review by Gibbon and Frith, caffeine had inconsistent effects on orthostatic hypotension, but no serious adverse events were reported⁵.

3.11 Sitting up

Night positioning

Elevating the head of the bed at night by 10-20° could decrease nocturnal hypertension and diuresis³.

Day Positioning

During the day, adequate orthostatic stress, ie, upright activity, should be maintained. If patients are repeatedly tilted up, their orthostatic hypotension is gradually attenuated, presumably by increasing venomotor tone.³

3.12 Exercise

Physical countermaneuvers can be performed to reduce *venous capacitance*, resulting in increased total peripheral resistance which assists venous return to the heart³.

Dosage

- Contracting the muscles below the waist for about 30 seconds at a time³

Examples

- Toe-raising³
- Leg-crossing and contraction³
- Thigh muscle co-contraction³
- Bending at the waist³
- Slow marching in place³
- SLR³

3.13 Patient Education

Considered the “single most important factor” in orthostatic hypotension management by Figueroa³.

Items to consider:

1. The mechanisms that maintain postural normotension and how to recognize the onset of orthostatic symptoms³.
2. There is no specific treatment of the underlying cause and that drug treatment alone is not adequate³.
3. Nonpharmacologic approaches and be aware that other drugs they start may worsen symptoms³.

Educate the patient on environmental stressors

- Prolonged or motionless standing
- Alcohol ingestion (causing vasodilation)
- Carbohydrate-heavy meals (causing postprandial orthostatic hypotension related to an increase in the splanchnic-mesenteric venous capacitance),
- Nocturnal diuresis causing early morning orthostatic hypotension
- Physical activity sufficient to cause muscle vasodilation
- Heat exposure (eg, hot weather or a hot bath or shower) producing skin vessel vasodilation
- Sudden postural changes
- Prolonged recumbency

3.14 Inpatient approach

- Perform motor and sensory evaluations first
- Move patient to sitting EOB as soon as possible
- Continue with subjective and objective
- (+) Hypotension
 - Have the patient drink water
 - Perform exercises sitting EOB

4 Dropfoot

Drop foot refers to a sign of motor weakness caused by [common fibular nerve palsy](#).

4.1 Etiology

During surgery, the nerve can be damaged through:

- Direct trauma⁶
- Thermal injury⁶
- Retractor placement⁶
- Hardware dislocation⁶
- Perforation⁶
- Postoperative Hematoma⁶
- Postoperative pseudotumor⁶

i Note

Up to 50% of the cases are idiopathic⁶

4.1.1 THA

Injury to [Common fibular division](#) of the [Sciatic nerve](#)⁶.

4.1.2 TKA

During a TKA, either the [common fibular division of the sciatic nerve](#) or the [Common fibular nerve](#) itself is damaged at some point during the operation.

4.2 Epidemiology

4.2.1 THA

This injury has been reported in 0.08% to 3.7% of primary arthroplasties and up to 7.6% in secondary or revision cases⁶.

4.2.2 TKA

Peroneal nerve palsy occurs in 0% to 9.5% of TKAs⁷.

4.3 Common fibular nerve vs Tibial Nerve

- Injuries to the sciatic nerve during total joint arthroplasty can affect the [Common fibular division](#) and/or the [tibial division](#) of the sciatic nerve⁶.
- Injuries to the [tibial division](#) are less severe and less common⁶.

4.4 Patient Presentation

4.4.1 Sensory symptoms

Sensory on dorsal aspect of the foot:

- Decreased sensation
- Numbness
- Tingling

4.4.2 Motor Symptoms

Loss of function in

- [Tibialis anterior](#)
- [Extensor digitorum longus](#)
- [Extensor hallucis longus](#)
- [Fibularis longus](#)
- [Fibularis brevis](#)
- [Fibularis tertius](#)

Resulting in

- Dorsiflexion weakness/paralysis ([footdrop](#))
- Eversion weakness/paralysis

4.4.3 Onset

Symptoms should be present by shortly after the operation in the recovery room or at the ward⁷.

4.5 Management

4.5.1 Patient education

If the [foot drop](#) is caused by the surgery, you should educate the patient that they need to advocate for themselves to home health physical therapy and outpatient physical therapy, that this issue was caused during the surgery and should be part of the physical therapy management.

4.5.2 Positioning

Passively positioning the ankle in dorsiflexion and eversion is important to prevent contractures⁸.

4.5.3 Stretching

Stretching the [Triceps surae](#) and the associated achilles tendon is important to prevent contractures⁸.

5 Anemia

5.1 Definition

Hemoglobin (HB) level is below the normative value:

- <13 g/dl for Males⁹
- <12 g/dl for Females⁹

5.2 Secondary complications

- acute kidney injury (AKI)⁹
- Delirium due to decreased oxygen transport⁹
 - Delirium often occurs postoperatively with an incidence of up to 74%⁹
- [Orthostatic intolerance](#)¹

5.3 Management

There is very little physical therapists can perform in the short term to prevent acute iatrogenic anemia.

At the same time, acute anemia will have little effect on whether a physical therapist can perform their evaluation and discharge, except for [orthostatic hypotension](#). Thus the best way to manage acute iatrogenic anemia, is to treat the symptoms of [OI](#) as outlined above¹.

References

1. Kurkis GM, Dennis DA, Johnson RM, Mejia M, Yazdani-Farsad Y, Jennings JM. Incidence and Risk Factors of Orthostasis After Primary Hip and Knee Arthroplasty. *The Journal of Arthroplasty*. 2022;37(6S):S70-S75. doi:[10.1016/j.arth.2022.01.035](https://doi.org/10.1016/j.arth.2022.01.035)
2. Mills PB, Fung CK, Travlos A, Krassioukov A. Nonpharmacologic management of orthostatic hypotension: A systematic review. *Archives of Physical Medicine and Rehabilitation*. 2015;96(2):366-375.e6. doi:[10.1016/j.apmr.2014.09.028](https://doi.org/10.1016/j.apmr.2014.09.028)
3. Figueroa JJ, Basford JR, Low PA. Preventing and treating orthostatic hypotension: As easy as A, B, C. *Cleveland Clinic Journal of Medicine*. 2010;77(5):298-306. doi:[10.3949/ccjm.77a.09118](https://doi.org/10.3949/ccjm.77a.09118)
4. Miller RH, Lowry JL, Meardon SA, Gillette JC. Lower extremity mechanics of iliotibial band syndrome during an exhaustive run. *Gait & Posture*. 2007;26(3):407-413. doi:[10.1016/j.gaitpost.2006.10.007](https://doi.org/10.1016/j.gaitpost.2006.10.007)
5. Gibbon JR, Frith J. The effects of caffeine in adults with neurogenic orthostatic hypotension: A systematic review. *Clinical Autonomic Research: Official Journal of the Clinical Autonomic Research Society*. 2021;31(4):499-509. doi:[10.1007/s10286-021-00814-5](https://doi.org/10.1007/s10286-021-00814-5)
6. Wu KY, Amrami KK, Hayford KM, Spinner RJ. Characterizing peroneal nerve injury clinicoradiological patterns with MRI in patients with sciatic neuropathy and foot drop after total hip replacement. *Journal of Neurosurgery*. 2023;139(6):1560-1567. doi:[10.3171/2023.5.JNS23173](https://doi.org/10.3171/2023.5.JNS23173)
7. Schinsky MF, Macaulay W, Parks ML, Kiernan H, Nercessian OA. Nerve injury after primary total knee arthroplasty. *The Journal of Arthroplasty*. 2001;16(8):1048-1054. doi:[10.1054/arth.2001.26591](https://doi.org/10.1054/arth.2001.26591)
8. Mohani MR, Arya N, Ratnani G, Harjpal P, Phansopkar P. Comprehensive Rehabilitation of a Patient With Foot Drop Secondary to Lumbar Canal Stenosis: A Case Report. *Cureus*. 2024;16(1):e52275. doi:[10.7759/cureus.52275](https://doi.org/10.7759/cureus.52275)
9. Kunz JV, Spies CD, Bichmann A, Sieg M, Mueller A. Postoperative anaemia might be a risk factor for postoperative delirium and prolonged hospital stay: A secondary analysis of a prospective cohort study. *PloS One*. 2020;15(2):e0229325. doi:[10.1371/journal.pone.0229325](https://doi.org/10.1371/journal.pone.0229325)